

CT CONTRAST QUESTIONNAIRE

Iodinated Intravenous contrast agents used for radiologic procedures, possess a small risk of reactions, which can range in severity from minor to fatal. Complications can be significantly reduced when we properly assess the patient's health status, obtain the medical history and review previous contrast reactions.

1. Previous **severe** contrast reaction: **STOP EXAM** __ Yes __ No
Anaphylaxis, convulsions cardiopulmonary arrest
2. Previous **moderate** contrast reaction: __ Yes __ No
Tachycardia, hypertension, hypotension, dyspnea, wheezing,
Laryngospasm, bronchospasm
3. Previous **mild** contrast reaction: __ Yes __ No
Nausea, vomiting, diaphoresis, itching, hives and/or rash
4. Are you allergic to Iodine? __ Yes __ No
5. History of allergic reaction to any substance. __ Yes __ No
6. History of asthma. __ Yes __ No
7. History of hypertension or hypotension. __ Yes __ No
8. History of kidney disease or dysfunction/renal disease. __ Yes __ No
9. History of diabetes: __ Yes __ No
Glucophage and Metformin-must be held for 48hrs following CT exam
10. History of sickle cell disease/multiple myeloma. __ Yes __ No
11. Are you pregnant or nursing an infant? __ Yes __ No
12. Have you eaten or had fluids in the last 4 hours? __ Yes __ No
13. Have you had a barium study in the last week? __ Yes __ No

I have been informed and understand the risks involved with contrast injection.

I, _____, consent to proceed with the following procedure: _____.

Patient Signature

Date

Witness

Date